



US008193381B2

(12) **United States Patent**  
**Lilga et al.**

(10) **Patent No.:** **US 8,193,381 B2**  
(45) **Date of Patent:** **Jun. 5, 2012**

(54) **HYDROXYMETHYL FURFURAL OXIDATION METHODS**

(75) Inventors: **Michael A. Lilga**, Richland, WA (US);  
**Richard T. Hallen**, Richland, WA (US);  
**Jianli Hu**, Overland Park, KS (US);  
**James F. White**, Richland, WA (US);  
**Michel J. Gray**, Pasco, WA (US)

(73) Assignee: **Battelle Memorial Institute**, Richland, WA (US)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 68 days.

(21) Appl. No.: **12/712,095**

(22) Filed: **Feb. 24, 2010**

(65) **Prior Publication Data**

US 2010/0152469 A1 Jun. 17, 2010

**Related U.S. Application Data**

(62) Division of application No. 11/932,436, filed on Oct. 31, 2007, now Pat. No. 7,700,788.

(60) Provisional application No. 60/863,704, filed on Oct. 31, 2006.

(51) **Int. Cl.**

**C07D 307/02** (2006.01)

**C07D 307/48** (2006.01)

(52) **U.S. Cl.** ..... **549/485**; 549/489

(58) **Field of Classification Search** ..... 549/483,  
549/484, 485, 488, 489  
See application file for complete search history.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

3,326,944 A \* 6/1967 Lew ..... 549/485  
4,549,025 A 10/1985 Dalcanele et al.  
4,897,497 A 1/1990 Fitzpatrick

4,977,283 A \* 12/1990 Leupold et al. .... 549/484  
5,608,105 A 3/1997 Fitzpatrick  
5,892,107 A 4/1999 Farone et al.  
6,706,900 B2 3/2004 Grushin et al.  
6,790,997 B2 9/2004 Eckert et al.  
7,956,203 B2 6/2011 Grushin et al.  
2003/0055271 A1 3/2003 Grushin et al.

**FOREIGN PATENT DOCUMENTS**

DE 3826073 A1 7/1988  
EP 0356703 A 3/1990  
FR 2669636 A1 5/1992  
GB 2188927 A 10/1987

(Continued)

**OTHER PUBLICATIONS**

Verdeguer et al, Catalytic Oxidation of HMF into furan-2,5-dicarboxylic acid, Journal of Molecular Catalysis, 1993, p. 327-344. translated.\*

(Continued)

*Primary Examiner* — Taylor Victor Oh

(74) *Attorney, Agent, or Firm* — Wells St. John P.S.

(57)

**ABSTRACT**

A method of oxidizing hydroxymethylfurfural (HMF) includes providing a starting material which includes HMF in a solvent comprising water into a reactor. At least one of air and O<sub>2</sub> is provided into the reactor. The starting material is contacted with the catalyst comprising Pt on a support material where the contacting is conducted at a reactor temperature of from about 50° C. to about 200° C. A method of producing an oxidation catalyst where ZrO<sub>2</sub> is provided and is calcined. The ZrO<sub>2</sub> is mixed with platinum (II) acetylacetonate to form a mixture. The mixture is subjected to rotary evaporation to form a product. The product is calcined and reduced under hydrogen to form an activated product. The activated product is passivated under a flow of 2% O<sub>2</sub>.

**15 Claims, 39 Drawing Sheets**

